

Surgical Associates of Austin, P.A.

F. Ames Smith, Jr., M.D., F.A.C.S.
F. William Cherico, Jr., M.D., F.A.C.S.
Robert M. Markus, Jr., M.D., F.A.C.S.
William D. Mayer, M.D., F.A.C.S.
R.Y. Declan Fleming, M.D., F.A.C.S.
Rodney B. Ashworth, M.D., F.A.C.S.
Peter V. Ching, M.D.
Aravind B. Sankar, M.D., F.A.C.S.
Alejandro Esquivel, M.D.

POST OPERATIVE INSTRUCTIONS

- Your incision will be closed with dissolvable sutures in most cases. There are some exceptions to this in which sutures would be placed for removal at a later office visit. You will be informed if this exception occurs. A sterile dressing will be placed over the incision to cover and should be **LEFT IN PLACE FOR THREE DAYS**. Upon removing the dressing, you will notice tapes across the incision; these should be left in place until they fall off. Also in some cases you will not have a dressing Derma bond will be used and in this case please leave glue on as long as possible. You may **SHOWER ON THE DAY FOLLOWING SURGERY**, but do not immerse the incision.
- Some degree of swelling, bruising, or numbness is expected at the operative site. You may place an ice pack at the site for 24-48 hours. You do not need to report these symptoms, as they will typically improve gradually without any special treatment.
- Please report any fever > (greater than) 101, bleeding that soaks through the dressing, or local signs of infection such as redness or drainage from the incision.
- Avoid stress at the suture line such as lifting, pulling, or pushing. For hernia surgery and any abdominal surgery, **PLEASE AVOID LIFTING** > (greater than) 20 pounds until approved by your doctor (typically this would be for at least 2 weeks).
- Please call the office (within 24-48 hours after going home) and make a follow-up appointment for about 2 weeks after the surgery.
- Generally, you should plan to eat lightly after surgery (liquids, soups, etc.) until any nausea related to anesthesia has lessened. This would usually apply to the evening after surgery, but after general anesthesia it might involve the next day as well.
- Avoid constipation by drinking plenty of fluids, taking a laxative as needed, and/or taking a psyllium product such as Metamucil or FiberCon. The pain medicines frequently can be associated with constipation.
- No driving for 3-4 days after surgery (it is ok to ride in the car). Avoid alcohol for at least 24 hours after surgery, or for as long as you are taking the pain medication.
- Call one of our nurses if you have other concerns or do not understand these instructions.
 - North office Monday & Tuesday 512-836-3230
 - **Central office--Elizabeth 512-472-1381 ext 112**
 - South office--Christina-512-447-4993 ext 202

Central Office
Plaza St. David
1015 E. 32nd, Suite 308
Austin, Texas 78705
512/472-1381
Fax 512/472-9688

North Office
Medical Oaks Pavilion
12201 Renfert Way
Suite 335
Austin, Texas 78758
512/836-3210
Fax 512/339-8203

South Office
South Austin Professional Building
4007 James Casey Street
Suite A-220
Austin, Texas 78745
512/447-4993
Fax 512/447-7084